

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number
WO 2006/068990 A1

(51) International Patent Classification:
C07D 205/08 (2006.01)

(21) International Application Number:
PCT/US2005/045901

(22) International Filing Date:
19 December 2005 (19.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/637,594 20 December 2004 (20.12.2004) US

(71) Applicant (for all designated States except US): **SCHERING CORPORATION** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033-0530 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **THIRUVETTIPURAM, Thiruvengadam, K.** [US/US]; 17 Village Road, Kendall Park, New Jersey 08824 (US). **CHIU, John, S.** [US/US]; 12 Ferndale Drive, Parsippany, New Jersey 07054 (US). **FU, Xiaoyong** [US/US]; 18 Augusta Avenue, Edison, New Jersey 08820 (US). **MCALLISTER, Timothy, L.** [US/US]; 45-A-2 Sandra Circle, Westfield, New Jersey 07090 (US).

(74) Agent: **KALYANARAMAN, Palaiyur, S.**; Schering-Plough Corporation, 2000 Galloping Hill Road, Mail Stop K-6-1 1990, Kenilworth, New Jersey 07033-0530 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE SYNTHESIS OF AZETIDINONES

(57) Abstract: A process is provided for preparing azetidinones useful as intermediates in the synthesis of penems and as hypocholesterolemic agents, comprising reacting a β -(substituted-amino)amide, a β -(substituted-amino)acid ester, or a β -(substituted-amino)thiolcarboxylic acid ester with a silylating agent and a cyclizing agent selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof, or the reaction product of: (i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or (ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride, wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl.



WO 2006/068990 A1

PROCESS FOR THE SYNTHESIS OF AZETIDINONES

BACKGROUND OF THE INVENTION


Field of the Invention

The present invention relates to a process for producing azetidinone or β -lactam compounds useful for treating vascular and lipidemic conditions, as well as intermediates for the synthesis of penems.

Azetidinones are useful as hypocholesterolemic agents, as disclosed in US RE 37,721, and also are useful as intermediates in the synthesis of penems, a known group of antibacterials.


EP Patent No. 0 707 567 discloses processes for preparing azetidinones by reacting a silylating agent and a fluoride ion catalyst cyclizing agent with a suitably protected β -(substituted-amino)amide, a β -(substituted-amino)acid ester, or a β -(substituted-amino)thiolcarbonic acid ester.

US Patent No. 6,207,822 discloses a process for making the compound (ezetimibe):



comprising:

(a) reacting p-fluorobenzoylbutyric acid with pivaloyl chloride and acylating the product with a chiral auxiliary to obtain a ketone:



wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is C₁-C₆ alkyl, phenyl, naphthyl, substituted phenyl, substituted naphthyl, C₁-C₆ alkoxy carbonyl or


- 2 -

benzyl, wherein the substituents on phenyl and naphthyl are 1-3 substituents selected from the group consisting of C₁-C₆ alkyl, phenyl and benzyl;

(b) reducing the ketone in the presence of a chiral catalyst to an alcohol:

(c) reacting the chiral alcohol of step (b), an imine and a silyl protecting agent,

5 then condensing the protected compounds to obtain a β -(substituted-amino)amide:



(d) cyclizing the β -(substituted-amino)amide with


(i) a silylating agent and a fluoride ion catalyst cyclizing agent;

(ii) a silylating agent and a quaternary ammonium salt of a chiral auxiliary of step (a);

10 or

(iii) a strong non-nucleophilic base;

to obtain the compound:



and removing the protecting groups.

15

Despite the above valuable improvements in preparing azetidinones, there remains a need for alternative, simpler and less expensive processes for preparing azetidinone and β -lactam compounds.

- 3 -

SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a process for preparing an azetidinone, or pharmaceutically acceptable salts, solvates, and prodrugs thereof,
5 comprising the step of reacting:

(1) a β -(substituted-amino)amide, a β -(substituted-amino)-acid ester, or a β -(substituted-amino)thiolcarbonic acid ester with

(2) at least one silylating agent and

(3) at least one cyclizing agent which is

(a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof, or


(b) the reaction product of:

(i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or

(ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride,

wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl.


Another aspect of the present invention is a process for preparing a compound
25 represented by the Formula (IV):



comprising the steps of:

cyclizing the β -(substituted-amino)amide of formula XII

- 4 -



wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxy carbonyl, and wherein Prot is a silyl protecting group

with at least one silylating agent and at least one cyclizing agent which is

(a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof, or


(b) the reaction product of:

(i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or

(ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride,

wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl,

to obtain the compound of Formula XI:



XI; and

removing the silyl protecting groups to form the compound of Formula (IV).

- 5 -


Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about."

DETAILED DESCRIPTION

In its many embodiments, the present invention provides a novel process for preparing azetidinones or β -lactams which can be useful as hypocholesterolemic or as intermediates in the synthesis of penems, a known group of antibacterials.

This process is applicable for preparing azetidinones which are optionally mono-, di- or unsubstituted at each of the C-3 and C-4 positions and substituted at the ring nitrogen. The stereochemistry of C-3, C-4-disubstituted azetidinones prepared by this process is dependent on the starting material: racemic, stereospecific or enantiomeric compounds can be obtained when the corresponding starting materials are used. In particular, this process is useful for the stereospecific preparation of azetidinones substituted in the C-3 and C-4 positions, and optionally substituted at the ring nitrogen.

The processes of the present invention can be used to prepare azetidinones such as are represented by the structural formula (I):



or pharmaceutically acceptable salts or solvates of the azetidinone of Formula (I), wherein in Formula (I) above:

X', Y' and Z can be the same or different and each is independently selected from the group consisting of $-\text{CH}_2-$, $-\text{CH}(\text{alkyl})-$ and $-\text{C}(\text{alkyl})_2-$;

- 6 -

Q¹ and Q² can be the same or different and each is independently selected from the group consisting of H, -(C₀-C₃₀ alkylene)-G, -OR⁶, -OC(O)R⁶, -OC(O)OR⁹, and -OC(O)NR⁶R⁷M;

Q³ is 1 to 5 substituents independently selected from the group consisting of
 5 acyl, alkyl, alkylaryl, alkylheteroaryl, alkylsulfonyl alkenyl, alkoxy, alkoxy carbonyl, alkynyl, -(C₀-C₃₀ alkylene)-G,
 -(C₀-C₁₀ alkylene)-OR⁶, -(C₀-C₁₀ alkylene)-C(O)R⁶,
 -(C₀-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-OC(O)R⁶,
 -(C₀-C₁₀ alkylene)-OC(O)OR⁹, -CH=CH-C(O)R⁶, -CH=CH-C(O)OR⁶,
 10 -C≡C-C(O)OR⁶, -C≡C-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-OR⁶,
 -O-(C₁-C₁₀ alkylene)-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-C(O)OR⁶, -CN,
 -C(=N-CN)-NH₂, -C(=NH)-NHR¹⁰, -O-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -O-(C₀-C₁₀ alkylene)-C(O)NR⁶NR⁷C(O)OR⁶,
 -O-(C₁-C₁₀ alkylene)-C(O)(aryl)-N-N=N⁻,
 15 -OC(O)-(C₁-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -(C₀-C₁₀ alkylene)-OC(O)NR⁶R⁷, -NO₂, -(C₀-C₁₀ alkylene)-NR⁶R⁷,
 -O-(C₂-C₁₀ alkylene)-NR⁶R⁷, -NR⁶C(O)R⁷, -NR⁶C(O)OR⁹,
 -NR⁶C(O)NR⁷R⁸, -NR⁶S(O)₀₋₂R⁹, -N(S(O)₀₋₂R⁹)₂, -CHNOR⁶, -C(O)NR⁶R⁷,
 -C(O)NR⁶NR⁶R⁷, -S(O)₀₋₂NR⁶R⁷, -S(O)₀₋₂R⁹,
 20 -O-C(O)-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -OC(O)-(C₁-C₁₀ alkylene)-NR⁶C(O)O-(alkylaryl), -P(O)(OR¹⁰)₂,
 -(C₁-C₁₀ alkylene)-OSi(alkyl)₃, -CF₃, -OCF₃, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,
 alkoxy carbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidioyl, allyl, allyloxy,
 aryloxy carbonyl, aralkoxy carbonyl (arylalkoxy carbonyl), aryl, arylalkyl (aralkyl),
 25 aryloxy, arylsulfonyl arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, benzoylbenzoyloxy,
 carboxy, cyano, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl,
 heteroarylalkenyl, heteroarylalkynyl, heteroarylalkyl, heteroarylalkoxy,
 heteroarylsulfonyl, heteroarylthio, dioxolanyl, heterocyclyl, heterocyclylalkyl,
 heterocyclylcarbonyl, heterocyclylcarbonylalkoxy, hydroxy, hydroxyalkyl, alkylsulfonyl,

- 7 -

and ring substituents comprising a moiety replacing one available hydrogen atom on a ring system or simultaneously replacing two available hydrogens on adjacent carbon atoms on a ring system as defined below;

Q⁴ is 1 to 5 substituents independently selected from the group consisting of

5 acyl, alkyl, alkylaryl, alkylheteroaryl, alkylsulfonyl alkenyl, alkoxy, alkoxy carbonyl, alkynyl,, -(C₀-C₃₀ alkylene)-G,

-(C₀-C₁₀ alkylene)-OR⁶, -(C₀-C₁₀ alkylene)-C(O)R⁶,

-(C₀-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-OC(O)R⁶,

-(C₀-C₁₀ alkylene)-OC(O)OR⁹, -CH=CH-C(O)R⁶, -CH=CH-C(O)OR⁶,

10 -C≡C-C(O)OR⁶, -C≡C-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-OR⁶,

-O-(C₁-C₁₀ alkylene)-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-C(O)OR⁶, -CN,

-C(=N-CN)-NH₂, -C(=NH)-NHR¹⁰, -O-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,

-O-(C₀-C₁₀ alkylene)-C(O)NR⁶NR⁷C(O)OR⁶,

-O-(C₁-C₁₀ alkylene)-C(O)(aryl)-N=N=N⁻,

15 -OC(O)-(C₁-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-C(O)NR⁶R⁷,

-(C₀-C₁₀ alkylene)-OC(O)NR⁶R⁷, -NO₂, -(C₀-C₁₀ alkylene)-NR⁶R⁷,

-O-(C₂-C₁₀ alkylene)-NR⁶R⁷, -NR⁶C(O)R⁷, -NR⁶C(O)OR⁹, -NR⁶C(O)NR⁷R⁸,

-NR⁶S(O)₀₋₂R⁹, -N(S(O)₀₋₂R⁹)₂, -CHNOR⁶, -C(O)NR⁶R⁷, -C(O)NR⁶NR⁶R⁷,

-S(O)₀₋₂NR⁶R⁷, -S(O)₀₋₂R⁹, -O-C(O)-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,

20 -OC(O)-(C₁-C₁₀ alkylene)-NR⁶C(O)O-(alkylaryl), -P(O)(OR¹⁰)₂,

-(C₁-C₁₀ alkylene)-OSi(alkyl)₃, -CF₃, -OCF₃, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxy carbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidioyl, allyl, allyloxy, aryloxy carbonyl, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, arylsulfonyl, aroylaroyloxy, aroyl, arylalkoxy carbonyl, benzoylbenzoyloxy, carboxy, cyano,

25 cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl, heteroarylalkenyl, heteroarylalkynyl, heteroarylalkyl, heteroarylalkoxy, heteroarylsulfonyl, heteroarylthio, dioxolanyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy, hydroxy, hydroxyalkyl, alkylsulfonyl, and ring substituents comprising a moiety replacing one available hydrogen atom on a

- 8 -

ring system or simultaneously replacing two available hydrogens on adjacent carbon atoms on a ring system as defined below;

Q^5 is 1 to 5 substituents independently selected from the group consisting of acyl, alkyl, alkylaryl, alkylheteroaryl, alkylsulfonyl alkenyl, alkoxy, alkoxy carbonyl, alkynyl, $-(C_0-C_{30} \text{ alkylene})-G$,

$-(C_0-C_{10} \text{ alkylene})-OR^6$, $-(C_0-C_{10} \text{ alkylene})-C(O)R^6$,
 $-(C_0-C_{10} \text{ alkylene})-C(O)OR^6$, $-(C_0-C_{10} \text{ alkylene})-OC(O)R^6$,
 $-(C_0-C_{10} \text{ alkylene})-OC(O)OR^9$, $-CH=CH-C(O)R^6$, $-CH=CH-C(O)OR^6$,
 $-C\equiv C-C(O)OR^6$, $-C\equiv C-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-OR^6$,

$-O-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-CN$,
 $-C(=N-CN)-NH_2$, $-C(=NH)-NHR^{10}$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,
 $-O-(C_0-C_{10} \text{ alkylene})-C(O)NR^6NR^7C(O)OR^6$,
 $-O-(C_1-C_{10} \text{ alkylene})-C(O)(\text{aryl})-N=N=N^+$,
 $-OC(O)-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$,

$-(C_0-C_{10} \text{ alkylene})-C(O)NR^6R^7$, $-(C_0-C_{10} \text{ alkylene})-OC(O)NR^6R^7$, $-NO_2$,
 $-(C_0-C_{10} \text{ alkylene})-NR^6R^7$, $-O-(C_2-C_{10} \text{ alkylene})-NR^6R^7$, $-NR^6C(O)R^7$,
 $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_{0-2}R^9$, $-N(S(O)_{0-2}R^9)_2$,
 $-CHNOR^6$, $-C(O)NR^6R^7$, $-C(O)NR^6NR^6R^7$, $-S(O)_{0-2}NR^6R^7$, $-S(O)_{0-2}R^9$,
 $-O-C(O)-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,

$-OC(O)-(C_1-C_{10} \text{ alkylene})-NR^6C(O)O-(\text{alkylaryl})$, $-P(O)(OR^{10})_2$,
 $-(C_1-C_{10} \text{ alkylene})-OSi(\text{alkyl})_3$, $-CF_3$, $-OCF_3$, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,
alkoxycarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidoyl, allyl, allyloxy,
aryloxycarbonyl, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, arylsulfonyl,
aroylaroyloxy, aroyl, arylalkoxycarbonyl, benzoylbenzoyloxy, carboxy, cyano,
cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl,
heteroarylalkenyl, heteroarylalkynyl, heteroarylalkyl, heteroarylalkoxy,
heteroarylsulfonyl, heteroarylthio, dioxolanyl, heterocyclyl, heterocyclylalkyl,
heterocyclylcarbonyl, heterocyclylcarbonylalkoxy, hydroxy, hydroxyalkyl, alkylsulfonyl,
and ring substituents comprising a moiety replacing one available hydrogen atom on a

- 9 -

ring system or simultaneously replacing two available hydrogens on adjacent carbon atoms on a ring system as defined below;

wherein optionally one or more carbon atoms of the $-(C_0-C_{30} \text{ alkylene})-$ radical of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is independently replaced by

5 $-O-$, $-C(O)-$, $-CH=CH-$, $-C\equiv C-$, $-N(\text{alkyl})-$, $-N(\text{alkylaryl})-$ or $-NH-$;

G is selected from the group consisting of a sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue, oligopeptide residue comprising 2 to 9 amino acids, trialkylammoniumalkyl radical and $-S(O)_2-OH$,

10 R^2 and R^3 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl and aryl;

R^6 , R^7 and R^8 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, and a protecting group; and

15 each R^9 is independently alkyl, cycloalkyl, aryl or arylalkyl.

each R^{10} is independently H or alkyl;

q is 0 or 1;

r is 0 or 1;

20 m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5.

As used above, and throughout the specification, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

25 The statements above, wherein, for example, Q^1 and Q^2 are said to be independently selected from a group of substituents, means that Q^1 and Q^2 are independently selected, but also that where an Q^1 or Q^2 variable occurs more than once in a molecule, those occurrences are independently selected (e.g., if Q^1 is $-OR^6$ wherein R^6 is hydrogen, Q^2 can be $-OR^6$ wherein R^6 is alkyl). Those skilled in the art will recognize that the size and nature of the substituent(s) will affect the number of
30 substituents that can be present.

- 10 -

The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties. It should be noted that any atom with unsatisfied valences in the text, schemes, examples and tables herein is assumed to have the hydrogen atom(s) to satisfy the valences.

5 The following definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Therefore, the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portions of "hydroxyalkyl", "haloalkyl", "alkoxy", etc.

10 As used herein, the term "acyl" means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

15 As used herein, the term "alkyl" means an aliphatic hydrocarbon group that can be straight or branched and comprises 1 to about 20 carbon atoms in the chain. Preferred alkyl groups comprise 1 to about 12 carbon atoms in the chain. More preferred alkyl groups comprise 1 to about 6 carbon atoms in the chain. "Branched" means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. The alkyl group can be substituted by one or more substituents independently selected from the group consisting of halo, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂ (which
20 alkyls can be the same or different), carboxy and -C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, heptyl, nonyl, decyl, fluoromethyl, trifluoromethyl and cyclopropylmethyl.

25 "Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as described herein. Preferred alkylaryls comprise a lower alkyl group. Non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl. "Alkylheteroaryl" means an alkylaryl moiety in which the aryl group comprises a heteroaryl group as defined herein.

30 "Alkylsulfonyl" means an alkyl-S(O₂)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

- 11 -

"Alkylsulfonyl" means an alkyl-S(O₂)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

"Arylsulfonyl" means an aryl-S(O₂)- group. The bond to the parent moiety is through the sulfonyl.

"Alkenyl" means an aliphatic hydrocarbon group (straight or branched carbon chain) comprising one or more double bonds in the chain and which can be conjugated or unconjugated. Useful alkenyl groups can comprise 2 to about 15 carbon atoms in the chain, preferably 2 to about 12 carbon atoms in the chain, and more preferably 2 to about 6 carbon atoms in the chain. The alkenyl group can be substituted by one or more substituents independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, heteroaryl, and alkoxy. When substituted the alkenyl group is described sometime herein for convenience by appending the substituent name, for example, when substituted with a heteroaryl group it can be described as a "heteroarylalkenyl" group. Non-limiting examples of suitable alkenyl groups include ethenyl, propenyl, n-butenyl, 3-methylbut-enyl and n-pentenyl.

Where an alkyl or alkenyl chain joins two other variables and is therefore bivalent, the terms alkylene and alkenylene, respectively, are used.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Useful alkoxy groups can comprise 1 to about 12 carbon atoms, preferably 1 to about 6 carbon atoms. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy and isopropoxy. The alkyl group of the alkoxy is linked to an adjacent moiety through the ether oxygen. Used successively, for example, alkoxyalkoxy, means an alkoxy moiety which is itself substituted with one or more alkoxy moieties.

"Alkoxyarylalkoxy" means an alkyl-O-aryl-alkylene-O- group in which the alkyl, alkylene and aryl groups are as previously described. Useful alkoxyarylalkoxy groups can comprise 7 to about 26 carbon atoms, preferably 7 to about 12 carbon atoms. A non-limiting example of a suitable alkoxyarylalkoxy group is methoxybenzyloxy. The alkoxyarylalkoxy is linked to an adjacent moiety through the ether oxygen.

- 12 -

"Alkoxy carbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxy carbonyl groups include methoxy carbonyl and ethoxy carbonyl. The bond to the parent moiety is through the carbonyl

"Aryloxy carbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxy carbonyl groups include phenoxy carbonyl and naphthoxy carbonyl. The bond to the parent moiety is through the carbonyl.

"Aralkoxy carbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxy carbonyl group is benzyloxy carbonyl. The bond to the parent moiety is through the carbonyl. For convenience the term "arylaralkoxy carbonyl" is sometimes used alternatively herein.

"Alkoxy carbonylalkoxy" means an alkyl-O-C(O)-alkylene-O- group in which the alkyl and alkylene groups are as previously described. Useful alkoxy carbonylalkoxy groups can comprise 3 to about 12 carbon atoms, preferably 3 to about 8 carbon atoms. A non-limiting example of a suitable alkoxy carbonylalkoxy group is CH₃CH₂-O-C(O)-CH₂-O-. The alkoxy carbonylalkoxy is linked to an adjacent moiety through the ether oxygen.

"Alkoxy iminoalkyl" means an alkyl-O-N=CH-alkylene- group in which the alkyl and alkylene groups are as previously described. Useful alkoxy iminoalkyl groups can comprise 2 to about 12 carbon atoms, preferably 2 to about 8 carbon atoms. The alkoxy iminoalkyl is linked to an adjacent moiety through the alkylene group.

"Alkyldioyl" means an ROC(O)-alkylene-C(O)-O- group in which R is alkyl or H and the alkylene group is as previously described. Useful alkyldioyl groups can comprise 2 to about 12 carbon atoms, preferably 2 to about 8 carbon atoms. Non-limiting examples of suitable alkyldioyl groups include 1,3-propanediol. The alkyldioyl is linked to an adjacent moiety through the ester oxygen.

"Alkynyl" means an aliphatic hydrocarbon group comprising at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butyne, 3-methylbutynyl, n-

- 13 -

pentynyl, and decynyl. The alkynyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, heteroaryl, and alkoxy. When substituted the alkynyl group is described sometime
5 herein for convenience by appending the substituent name, for example, when substituted with a heteroaryl group it can be described as a "heteroarylalkynyl" group.

"Allyl" means $(R^2R^3)C=CR^2-C(R^2R^3)-$, wherein R^2 and R^3 can be the same or different and are selected independently for each occurrence from the group consisting of hydrogen, alkyl and aryl;

10 "Allyloxy" means $H_2C=CH-O-$. The allyloxy is linked to an adjacent moiety through the ether oxygen.

"Aryl" means an aromatic monocyclic or multicyclic ring system comprising about 5 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The
15 aryl group can be substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl, naphthyl, indenyl, tetrahydronaphthyl and indanyl.

"Arylene" means a bivalent phenyl group, including ortho, meta and para-substitution.

"Aralkyl" or "arylalkyl" means an aryl-alkylene- group in which the aryl and
20 alkylene are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, phenethyl and naphthylmethyl. The aralkyl is linked to an adjacent moiety through the alkylene group.

"Aryloxy" means an aryl-O- group in which the aryl group is as previously
25 described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

"Aralkoxy" or "arylalkyloxy" means an aralkyl-O- group in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkoxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is
30 through the ether oxygen. "Aralkoxycarbonyl" means an aralkoxy-C(O)- group in which the aralkoxy group is as previously described.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1- and 2-naphthoyl.

"Aroyloxy" means an aroyl-O- group in which the aroyl group is as previously described. The bond to the parent moiety is through the ether oxygen. Non-limiting examples of suitable groups include benzoyloxy and 1- and 2-naphthoyloxy.

"Carboxy" means R'-C(O)O-, wherein R' is an aliphatic or aromatic hydrocarbon radical which may or may not be substituted.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be substituted with one or more "ring system substituents" which may be the same or different, and are as defined below. Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like. "Cycloalkylene" refers to a corresponding bivalent ring, wherein the points of attachment to other groups include all positional isomers.


"Cycloalkylalkyl" means a cycloalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkylalkyls include cyclohexylmethyl, adamantylmethyl and the like.

"Cycloalkenyl" means a non-aromatic mono or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms which contains at least one carbon-carbon double bond. Preferred cycloalkenyl rings contain about 5 to about 7 ring atoms. The cycloalkenyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkenyls include cyclopentenyl, cyclohexenyl, cyclohepta-1,3-dienyl, and the like. Non-limiting example of a suitable multicyclic cycloalkenyl is norbornenyl.

"Cycloalkenylalkyl" means a cycloalkenyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkenylalkyls include cyclopentenylmethyl, cyclohexenylmethyl and the like.

"Dioxolanyl" means.

- 15 -



"Halo" refers to fluorine, chlorine, bromine or iodine radicals. Preferred are fluoro, chloro or bromo, and more preferred are fluoro and chloro.

5 "Heteroaryl" means a monocyclic or multicyclic aromatic ring system of about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is/are atoms other than carbon, for example nitrogen, oxygen or sulfur. The heteroatom(s) interrupt a carbocyclic ring structure and have a sufficient number of delocalized pi electrons to provide aromatic character, provided that the rings do not contain adjacent oxygen and/or sulfur atoms. Preferred
10 heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is
15 present as a ring atom. A nitrogen atom of a heteroaryl can be oxidized to form the corresponding N-oxide. All regioisomers are contemplated, e.g., 2-pyridyl, 3-pyridyl and 4-pyridyl. Examples of useful 6-membered heteroaryl groups include pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl and the like and the N-oxides thereof. Examples of useful 5-membered heteroaryl rings include furyl, thienyl, pyrrolyl, thiazolyl,
20 isothiazolyl, imidazolyl, pyrazolyl and isoxazolyl. Useful bicyclic groups are benzo-fused ring systems derived from the heteroaryl groups named above, e.g., quinolyl, phthalazinyl, quinazolinyl, benzofuranyl, benzothienyl and indolyl.

"Heteroarylalkyl" or "heteroaralkyl" means a heteroaryl-alkylene- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain
25 a lower alkyl group. Non-limiting examples of suitable heteroaralkyl groups include pyridylmethyl, 2-(furan-3-yl)ethyl and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkylene. In the same manner, "heteroarylalkenyl" and "heteroarylalkynyl" mean a heteroaryl group, as defined herein, bonded to the parent moiety through, respectively, an alkenyl and an alkynyl group. "Heteroarylalkoxy"

- 16 -

means a heteroaryl-alkylene-O- group in which the heteroaryl and alkylene are as previously described.

"Heterocyclyl" means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,3-dioxolanyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like.

"Heterocyclylalkyl" means a heterocyclyl-alkylene- group in which the heterocyclyl and alkylene groups are as previously described. Preferred heterocyclylalkyls contain a lower alkylene group. The bond to the parent moiety is through the alkylene. "Heterocyclylcarbonyl" means a heterocyclyl-C(O)- group in which the heterocyclyl is as previously described.


Preferred heterocyclylcarbonyls contain a lower alkyl group. The bond to the parent moiety is through the carbonyl. "Heterocyclylcarbonylalkoxy" means a heterocyclyl-C(O)-alkoxy- group in which the heterocyclyl and alkoxy are as previously described.

"Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

"Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl,

- 17 -

heteroaryl, aralkyl, alkylaryl, aralkenyl, heteroaralkyl, heteroarylalkynyl, alkylheteroaryl, heteroaralkenyl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxycarbonyl, aryloxy carbonyl, aralkoxycarbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylsulfinyl, arylsulfinyl, heteroarylsulfinyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, , -C(=N-CN)-NH₂, -C(=NH)-NH₂, -C(=NH)-NH(alkyl), Y₁Y₂N-, Y₁Y₂N-alkyl-, Y₁Y₂NC(O)- Y₁Y₂NSO₂- and -SONY₁Y₂, wherein Y₁ and Y₂ may be the same or different and are independently selected from the group consisting of hydrogen, alkyl, aryl, and cycloalkyl, and aralkyl. "Ring system substituent" may also mean a single moiety which simultaneously replaces two available hydrogens on two adjacent carbon atoms (one H on each carbon) on a ring system. Examples of such moiety are methylene dioxy, ethylenedioxy, -C(CH₃)₂- and the like which form moieties such as, for example:



"Sugar residue" means a moiety derived from an aldose or ketose that has 3 to 7 carbon atoms and may belong to the D or L series. Non-limiting examples of suitable aldoses from which the sugar residue can be formed include glucose, mannose, galactose, ribose, erythrose and glyceraldehydes. A non-limiting example of a suitable ketose from which the sugar residue can be formed is fructose.

"Disugar residue" means a moiety derived from a sugar that can be hydrolyzed to two monosaccharide molecules. Non-limiting examples of suitable compounds from which the disugar residue can be formed include maltose, lactose, cellobiose and sucrose.

Examples of sugar residues and disugar residues include those moieties G listed in detail above.

Di-, tri- or tetrasaccharides are formed by acetal-like binding of two or more sugars. The bonds may be in α or β form. "Trisugar residue" means a moiety derived from a sugar that can be hydrolyzed to three monosaccharide molecules. "Tetrasugar

- 18 -

residue" means a moiety derived from a sugar that can be hydrolyzed to four monosaccharide molecules.

If the sugar is substituted, the substitution is preferably at the hydrogen atom of an OH group of the sugar.

5 "Sugar acid" means an sugar residue, such as can be formed from glucuronic acid, galacturonic acid, gluconic acid, galactonic acid, mannonic acid, glucaric acid and galactaric acid.

"Amino sugar" means an amino-substituted sugar residue such as can be formed from glucosamine, galactosamine, glucamine or 3-amino-1,2-propanediol.


10 Suitable protective groups for the hydroxyl groups of the sugars include benzyl, acetyl, benzoyl, pivaloyl, trityl, tert-butyldimethylsilyl, benzilidene, cyclohexidene or isopropylidene protective groups.

"Amino acid residue" means a moiety derived from an amino acid. The amino acid moiety can be prepared from the D or L forms of the amino acid. Non-limiting
15 examples of suitable amino acids from which the amino acid residue can be prepared include alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, histidine, hydroxylysine, hydroxyproline, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophane, tyrosine, valine, 2-aminoadipic acid, 3-aminoadipic acid, beta-alanine, 2-aminobutyric acid, 4-
20 aminobutyric acid, piperidino carboxylic acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-(2-thienyl)glycine, penicillamine, N-ethylasparagine, 2-aminoisobutyric acid, 2-aminoisobutyric acid, 2-aminopimelic acid, 2,4-diaminobutyric acid, desmosine, 2,2-diaminopimelic acid, 2,3-diaminopropioninc acid, N-ethylglycine, 3-(2-thienyl)alanine, sarcosine, N-methylisoleucine, 6-N-methyllysine, N-methylvaline, norvaline,
25 norleucine, ornithine and N-methylglycine.

"Oligopeptide residue" means the residue of a peptide constructed of 2 to 9 of the amino acids mentioned above.

"Trialkylammonium alkyl radical" means the group

- 19 -



wherein n_1 is 0 to 10 and Alk_1 , Alk_2 and Alk_3 can be the same or different and each is a straight or branched alkyl radical having 1 to 20 carbon atoms.

Azetidinone or β -lactam compounds prepared in accordance with the methods of the invention have at least one asymmetrical carbon atom. Accordingly, the methods of the invention are useful in the preparation of all isomers, including enantiomers, stereoisomers, rotamers, tautomers and racemates of the compounds of Formula (I) (where they exist) and are contemplated as being within the scope of products formed by the process of this invention. Accordingly, the invention includes the preparation of d and l isomers in both pure form and in admixture, including racemic mixtures. Utilizing the methods of the invention, isomers can be prepared using conventional techniques, either by reacting optically pure or optically enriched starting materials or by separating isomers of a compound of the Formula (I). As the term is used herein, isomers may also include geometric isomers, e.g., when a double bond is present. Preparation of polymorphous forms of the compounds of Formula (I), whether crystalline or amorphous, also are contemplated as being part of this invention.

Azetidinone or β -lactam compounds prepared in accordance with the methods of the invention having an amino group can form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salt is prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt. The free base form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium bicarbonate. The free base form differs from its respective salt form somewhat in certain physical properties, such as solubility in polar solvents, but the salt is otherwise equivalent to its respective free base forms for purposes of the invention.


- 20 -

Certain azetidinone or β -lactam compounds prepared in accordance with the methods of the invention are acidic (e.g., those compounds which possess a carboxyl group). These compounds form pharmaceutically acceptable salts with inorganic and organic bases. Examples of such salts are the sodium, potassium, calcium,
5 aluminum, gold and silver salts. Accordingly, the methods of the invention are useful in the provision of salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.

Azetidinone or β -lactam compounds prepared in accordance with the methods of the invention having a carboxylic acid group can form pharmaceutically acceptable
10 esters with an alcohol. Examples of suitable alcohols include methanol and ethanol. The methods of the present invention are also useful in the provision of such compounds.

Non-limiting examples of azetidinones that can be prepared using the
15 *processes of the present invention are disclosed in US Patents Nos. RE 37,721, 5,624,920, 5,656,624, 5,627,176, 5,633,246, 5,661,145, 5,688,785, 5,688,787, 5,688,990, 5,744,467, 5,756,470, 5,767,115, 5,846,966, each incorporated by reference herein.*

Preferably, the processes of the present invention can be used to prepare the
20 azetidinone (ezetimibe) represented by the structural formula (IV):



(IV).

Ezetimibe is commercially available in the US in ZETIA™ formulation and in combination with simvastatin in VYTORIN™ formulation from MSP Pharmaceuticals,
25 Inc.

- 21 -


In one embodiment, the present invention provides a process for preparing an azetidinone, comprising reacting: (1) a β -(substituted-amino)amide, a β -(substituted-amino)-acid ester, or a β -(substituted-amino)thiolcarbonic acid ester with (2) at least one silylating agent and (3) at least one cyclizing agent described in detail below.

As used herein, the terms β -(substituted-amino)amide, β -(substituted-amino)acid ester, and β -(substituted-amino)thiolcarbonic acid ester refer to β -aminoamides, β -aminoacid esters, and β -aminothiol-carbonic acid esters refer to secondary amines, that is, compounds wherein the nitrogen is joined to the β -carbon, to a hydrogen molecule, and to a non-hydrogen substituent.

Starting β -(substituted-amino)amides, β -(substituted-amino)acid esters and β -(substituted-amino)thiolcarbonic acid esters are known or can be prepared by one skilled in the art using known methods. For example, suitable methods for preparing β -aminoamide compounds of Formula II are disclosed in WO 93/02048, incorporated by reference herein.

Those skilled in the art will recognize that for cyclization to proceed as desired, -NH₂, -SH and -OH substituents present on the β -(substituted-amino)amide, a β -(substituted-amino)acid ester, or a β -(substituted-amino)thiolcarbonic acid ester starting material must be converted to groups which will not be silylated, either preferentially or in addition to silylation of the substituted-amino portion of the molecule. Suitable protecting groups well known in the art include for -NH₂: t-butyltrimethylsilyl, benzyl, benzoyl and t-butoxy-carbonyl; for -SH: triphenylmethyl; and for -OH: lower alkoxy, e.g., methoxy, benzyloxy and t-butyltrimethylsilyl.

In one embodiment, the β -(substituted-amino)amide comprises a carbamoyl portion B-C(O)-, wherein B is a deprotonated chiral auxiliary selected from the group consisting of




- 22 -

wherein X is -O-, -S- or -N(alkyl)-; Y is =O or =S; and R¹² and R¹³ are each independently selected from the group consisting of alkyl, aryl and alkoxy carbonyl, or wherein one of R¹² or R¹³ is as defined above and the other is hydrogen. Preferably, X and Y are each oxygen and R¹² is hydrogen and R¹³ is phenyl, benzyl or isopropyl.

5 The aryl group of R¹² or R¹³ can be independently selected from the group consisting of phenyl, naphthyl, benzyl, substituted phenyl, substituted naphthyl and substituted benzyl, wherein the substituents on the phenyl, naphthyl or benzyl are 1-3 substituents selected from the group consisting of alkyl, alkoxy, phenyl and benzyl.

In another embodiment, the β -(substituted-amino)amide comprises a
10 carbamoyl portion B-C(O)-, where B is (R¹⁴)(R¹⁵)N-, and R¹⁴ and R¹⁵ are each independently selected from the group consisting of alkyl, aryl, arylalkyl.




Preferably, the β -(substituted-amino)amide is represented by Formula (II):
15 wherein B, R², R³, R⁶, R⁷, R⁸, R⁹, R¹⁰, Q¹, Q², Q³, Q⁴, Q⁵, X', Y', Z, m, n, p, q, r are as defined above.

More preferably, Q⁴ is selected from the group consisting of H, 3-fluoro-, and 4-fluoro- and Q⁵ is selected from the group consisting of H, 4-methyl-, 2-fluoro-, and 4-fluoro-.

Most preferably, the β -(substituted-amino)amide is represented by Formula
20 (III):

- 23 -



(III).

In another embodiment of the process of the present invention, a β -(substituted-amino)acid ester can be used as a reactant. Preferably, the β -(substituted-amino)acid ester comprises a carboxylic acid ester portion $R^{14}-O-C(O)-$,
 5 wherein R^{14} is alkyl or aryl.

In another embodiment of the process of the present invention, a β -(substituted-amino)thiolcarmonic acid ester can be used as a reactant. Preferably, the β -(substituted-amino)thiolcarmonic acid ester comprises a thiolcarmonic acid ester portion $R^{14}-S-C(O)-$, wherein R^{14} is alkyl or aryl.

10 As discussed above, the process comprises reacting (1) the β -(substituted-amino)amide, a β -(substituted-amino)-acid ester, or a β -(substituted-amino)thiolcarmonic acid ester with (2) one or more silylating agents and (3) one or more cyclizing agents described below.

15 Preferably the silylating agent is a silylenol ether. Non-limiting examples of suitable silylenol ethers include bistrimethylsilylacetamide, N-methyl-O-trimethylsilylacetamide or isopropenyloxy trimethylsilane. The silylating agent can be added in up to a stoichiometric amount or excess.

In the process of the present invention, suitable cyclizing agent(s) are: (a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof, or (b) the reaction product of:

(i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or

- 24 -

(ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride, wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted (i.e., quaternary ammonium) or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl.

Non-limiting examples of suitable quaternary ammonium moieties include alkylammonium groups containing alkyl groups of 1-6 carbon atoms, e.g., tetra n-butylammonium. Examples of suitable arylalkyl-alkylammonium groups include benzyltriethyl-ammonium and benzyl-trimethylammonium and examples of aryl-alkyl-ammonium include phenyltriethylammonium and phenyltrimethyl-ammonium.

The alkali metal carboxylate can be a carboxylate salt of an alkali metal selected from the group consisting of lithium, sodium, potassium and cesium, preferably potassium.

In one embodiment, the alkali metal carboxylate or quaternary ammonium carboxylate can comprise a carboxylate moiety selected from the group consisting of formate, acetate, propionate, butyrate, valerate, caproate, caprylate, laurate, myristate, palmitate, stearate, oleate, linoleate, linolenate, cyclohexanecarboxylate, phenylacetate, benzoate and toluate, preferably acetate.

In another preferred embodiment, the quaternary ammonium carboxylate is tetra n-butylammonium acetate. In another preferred embodiment, the silylating agent is bistrimethylsilylacetamide and the cyclizing agent is tetra n-butylammonium acetate.

In yet another preferred embodiment, the quaternary ammonium hydroxide is tetra n-butylammonium hydroxide.

As discussed above, the cyclizing agent can be the reaction product of (i) at least one quaternary ammonium halide and at least one alkali metal carboxylate in a catalytic to stoichiometric excess amount. Suitable quaternary ammonium moieties are discussed above. The quaternary ammonium halide can comprise a halide moiety selected from the group consisting of fluoride, chloride, bromide and iodide, preferably fluoride. In a preferred embodiment, the alkali metal carboxylate is potassium acetate.

In another embodiment, the cyclizing agent can be the reaction product of (ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride in a catalytic to

- 25 -

stoichiometric excess amount. In a preferred embodiment, the cyclizing agent is the reaction product of tetra n-butylammonium bromide and potassium acetate. In another preferred embodiment, the cyclizing agent is the reaction product of tetra n-butylammonium bromide and cesium fluoride.

5 The reaction product (3)(b) can be formed in situ in the presence of reactants (1) and (2) or formed prior to reaction with reactants (1) and (2). Advantages of forming the reaction product (3)(b) in situ include that the anhydrous reactants are easier to handle and it may be less expensive to prepare the cyclizing agent from the reactants.

10 When a hydrated quaternary ammonium cyclizing agent is used, the reagent is added in a catalytic amount, i.e., about 1 to about 20 mole percent, preferably about 5 mole percent, and when an anhydrous quaternary ammonium cyclizing agent is used, it can be added in a catalytic up to a stoichiometric amount. When an alkali metal carboxylate is used, it is added in catalytic amount up to a stoichiometric amount
15 compared to the starting β -amino compound. If added to the reaction mixture after the silylation agent, the fluoride reagent is added directly to the reaction mixture resulting from silylation, and is reacted at about 0°C to 110°C, preferably about 20°C to 60°C, for about 0.5 to about 6 hours, preferably about 1 hour.


20 In an alternative embodiment, one or more fluoride ion catalyst cyclizing agents can be used in combination with the cyclizing agents described above. Non-limiting examples of suitable fluoride ion catalyst cyclizing agents include quaternary alkyl-, aryl-alkyl- or arylalkyl-alkylammonium fluoride salt or a hydrate thereof, or a mixture thereof, wherein alkyl-, aryl-alkyl- or arylalkyl-alkylammonium are as defined above, or
25 is an alkali metal fluoride salt or a hydrate thereof, such as cesium fluoride or potassium fluoride. When a fluoride ion catalyst cyclizing agent is used, the agent is added in an amount sufficient to replace the cyclizing agent discussed above on a 1:1 molar basis as desired. If added to the reaction mixture after the silylation agent, the fluoride reagent is added directly to the reaction mixture resulting from silylation, and
30 is reacted at about 0°C to 110°C, preferably about 20°C to 60°C, for about 0.5 to about 6 hours, preferably about 1 hour. When the silylation reagent, cyclization agent and the fluoride reagent are added simultaneously, the reaction is conducted under similar conditions.

- 26 -

The order of addition of the components of this process is not critical to the preparation of the azetidinone product. For example, the starting β -(substituted-amino)amide, β -(substituted-amino)acid ester, or β -(substituted-amino)thiolcarmonic acid ester can first be reacted with the silylating agent and then reacted with the cyclizing agent, or the starting compound can be added to a mixture of the silylating agent and the cyclizing agent.

Silylation is effected by reacting the starting material with a silyl-enol ether silylating agent such as bistrimethylsilyl acetamide (BSA), N-methyl-O-trimethylsilyl acetamide or iso-propenyloxy trimethylsilane, preferably BSA, in a suitable inert organic solvent at 0°C to 110°C, preferably at about 20°C to 90°C, and more preferably at ambient temperature (e.g., about 25°C). The reaction is preferably carried out in a dry, inert atmosphere, e.g., the solvent is dried, typically with molecular sieves, and the reaction is carried out under nitrogen. When the silylation and cyclization are done sequentially, i.e., the silylating agent is reacted with the starting material first, the silylation reaction can be allowed to continue for up to about two hours, but preferably the cyclization step is carried out immediately after silylation, or the silylating agent and the cyclizing agent are added simultaneously.

Another aspect of the present invention is a process for preparing an azetidinone represented by the structural formula (I):




or pharmaceutically acceptable salts or solvates of the azetidinone of Formula (I), wherein in Formula (I) above: R^2 , R^3 , R^6 , R^7 , R^8 , R^9 , R^{10} , Q^1 , Q^2 , Q^3 , Q^4 , Q^5 , X' , Y' , Z , m , n , p , q , r are as above,


comprising the step of reacting:

(a) a compound of Formula (II):

- 27 -



wherein R_2 , R_3 , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , X' , Y' , Z , m , n , p , q , r are as above, and B is a




wherein X is $-O-$, $-S-$ or $-N(C_1-C_6 \text{ alkyl})-$; Y is $=O$ or $=S$; and R^{12} and R^{13} are each independently selected from the group consisting of alkyl, aryl and alkoxy carbonyl, or wherein one of R^{12} or R^{13} is as defined above and the other is hydrogen, or B is $(R^{14})(R^{15})N-$, wherein R^{14} and R^{15} are each independently selected from the group consisting of alkyl and aryl;

with (b) at least one silylating agent and (c) at least one cyclizing agent discussed above.

In a preferred embodiment, the process described above is used for preparing an azetidinone represented by the structural Formula (IV):

- 28 -



(IV)


comprising reacting a β -(substituted-amino)amide of Formula (III) above with (b) at least one silylating agent and (c) at least one cyclizing agent as described above.

5

Another aspect of the present invention is a process for preparing a compound represented by the structural Formula (IV) above, comprising:

(a) reacting *p*-fluorobenzoylbutyric acid of Formula V with pivaloyl chloride and acylating the product with a chiral auxiliary of Formula VI to obtain a ketone of Formula VII:

10



wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxy carbonyl;

(b) reducing the ketone of Formula VII in the presence of a chiral catalyst to an alcohol of Formula VIII:


15




;

(c) reacting the chiral alcohol of Formula VIII, an imine of formula IX and a silyl protecting agent, then condensing the silyl-protected compounds to obtain a β -(substituted-amino)amide of Formula XII, wherein Prot is a silyl protecting group:

- 29 -




(d) cyclizing the β -(substituted-amino)amide of formula XII with a silylating agent and a cyclizing agent discussed above, to obtain the compound of Formula XI:



(e) removing the silyl protecting groups.


In another embodiment, the present invention provides a process for preparing a compound represented by the Formula (IV) above, comprising:

reacting a chiral alcohol of Formula VIII, an imine of Formula IX and a silyl protecting agent, then condensing the silyl-protected compounds to obtain a β -(substituted-amino)amide of Formula XII, wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxycarbonyl, and wherein Prot is a silyl protecting group:




cyclizing the β -(substituted-amino)amide of Formula XII with a silylating agent and a cyclizing agent discussed above, to obtain the compound of Formula XI:

- 30 -




removing the silyl protecting groups.


- 5 Another aspect of the present invention is a process for preparing a compound represented by the Formula (IV):



cyclizing the β -(substituted-amino)amide of formula XII








- 10 with a silylating agent and a cyclizing agent discussed above, to obtain the compound of Formula XI:



- 15 Preferred reaction conditions are shown in the following scheme:

Step (a):

- 31 -


Step (b):Step (c):Step (d):Step (e):

In the reaction scheme above, TEA is triethylamine, DMAP is 4-dimethyl-amino pyridine, DIPEA is diisopropylethylamine, BSA is bistrimethylsilyl acetamide, TBAOAc


- 32 -

is tetra n-butyl-ammonium acetate, t-BuOMe is t-butyl methyl ether and Prot is a silyl protecting group as defined above.

Starting materials of Formulae V and VI are known in the art, and the procedure of step (a) for reacting a compound of Formula V and a compound of
 5 Formula VI is known in the art. Preferably, the chiral auxiliary of Formula VI is exemplified by the formula:



wherein Y is =O, X is -O- and R¹ is phenyl, benzyl or C₁-C₆ alkyl. A



See the example below for typical reaction conditions.

10 Similarly, the procedure of step (b) for reducing a ketone to a hydroxy group using a borane reducing agent such as BH₃•S(CH₃)₂ in the presence of a chiral catalyst such as (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo(1,2-c)(1,3,2) oxaza-borolidine is known: See U.S. RE 37,721, U.S. 6,207,822 and U.S. 6,627,757, each incorporated by reference herein.

15 In step (c), the chiral alcohol, VIII, and the imine, IX, are protected with a suitable hydroxy-protecting group, preferably a silyl protecting group such as that derived from chlorotrimethylsilane (TMSCl) or t-butyldimethyl-silyl chloride (TBDMSCl). The alcohol (1 equivalent) and imine (preferably 1-3 equivalents) are added to an anhydrous solvent such as CH₂Cl₂, the reaction mixture is cooled to -10°
 20 to 15°C, a tertiary amine base such as DIPEA is added (preferably 2-4 equivalents), and sufficient silylating reagent to react with both the alcohol and the imine is added (e.g., 2-4 equivalents). After silylation is complete, the alcohol and imine are condensed by reacting at -20° to -35°C with at least 1 equivalent of a Lewis acid such as TiCl₄, in the presence of a tertiary amine base (preferably 1-3 equivalents) such as

- 33 -

DIPEA for 2-4 hours. The reaction is quenched, for example by treating with an acid such as glacial acetic acid followed by aqueous tartaric acid solution; the resultant product is extracted and crystallized using conventional procedures.

5 The azetidinone resulting from the processes of the present invention can be purified by appropriate standard procedures such as column chromatography or crystallization.


10 The term "suitable inert organic solvent" as used above means any organic solvent or combination of solvents that is unreactive in the reaction being conducted and is a solvent for the reactants. Typical suitable solvents are halogenated compounds such as dichloromethane; heterocyclic compounds such as tetrahydrofuran (THF); DMSO; dimethyl-formamide (DMF); acetonitrile; and carbocyclic aromatics such as toluene. Preferred are toluene, THF and dichloromethane.

15 The following examples illustrate the process of this invention. Although the examples are directed to C-3, C-4 disubstituted compounds and the stereochemistry of the reactants and intermediates are indicated in the various depicted structural formulas in the following examples, it is to be understood that the process of this invention is operative for azetidinones regardless of stereochemistry, and involves
20 merely the selection of reactants having the desired racemic or stereochemical configuration and the selection of reaction conditions which result in the desired configuration in the product.


EXAMPLES

25 Starting materials useful in the processes of the present invention can be made by the following procedures:

- 34 -

Preparation of 4-(4-fluorobenzoyl)butyric acid:


Charged 250 g of anhydrous AlCl_3 (1.87 moles) to a 2 L 3-neck round bottom flask, added 300 mL fluorobenzene (307.5 g; 3.2 moles) and cooled the mixture in an ice bath to 5°C. Added a hazy suspension of 100 g glutaric anhydride (0.86 mole) in 400 mL fluorobenzene (4.3 moles) through an addition funnel over a period of 45 min., and maintained the temperature below 12°C. The reaction mixture was warmed to ambient temperature gradually and agitated at r.t. for about 90 min.; checked for completion by NMR. Cooled the reaction mixture to 0 to 5°C, then added a cold aqueous solution (700 mL) of 1N HCl carefully to the mixture to destroy any unreacted AlCl_3 , keeping the temperature of the mixture below 20°C during the early part of the acid addition, and below 40°C for the rest of the time. Poured the entire mixture into a 2 L 1:1 mixture of water and ice (v/w) to precipitate out crude products, filtered the white suspension and washed well with water. Added the white residue to 3 L of aqueous saturated solution (~5%) of NaHCO_3 , heated the basic mixture on a steam bath for one hour and filtered the batch while hot through a thin pad of celite. Cooled the filtrate to r.t., added about 320 mL of concentrated HCl dropwise into the filtrate to pH 1 to crystallize out products, and agitated the white suspension in an ice bath for 30 min. Filtered the batch, washed the wet cake with ice cold water and dried in a vacuum oven at 50°C for 16 h to obtain 143.2 g of 4-(4-fluorobenzoyl)-butyric acid; m.p. 141 to 142°C, isolated yield: 79.3%.

Preparation of Imine:

Equipped a three necked 1L flask with a mechanical stirrer, thermometer and an addition funnel. Added 480 mL of isopropanol, 144 g (1.18 moles) of p-

- 35 -

hydroxybenzaldehyde (endothermic) and agitated the mixture while heating to a temperature of 50°C. Agitated the mixture at 50°C for 15 min (making sure all the material was in solution), then added 114 mL (1.2 moles) of p-fluoroaniline slowly via the addition funnel (exothermic reaction). After the addition was complete, agitated the thick slurry for 1 hr at 50°C, cool to r.t. and agitated for 30 min. Filtered the product, washed the cake with 150 mL of isopropanol, dried the wet cake in a draft oven at 50°C for 24 h or until dry to yield 222 g of the imine (88%); mp: 180-182°C.

Process Example:**Step (a):**


Equipped a 3-necked 500 mL round bottom flask with a thermometer, an addition funnel and a nitrogen inlet. Added p-fluoro-benzoylbutyric acid (20 g, 95.15 mmol), CH₂Cl₂ (100 mL) and TEA (23 mL, 165 mmol) and agitated the mixture at room temperature for 5 min. Added trimethylacetyl chloride (11.3 mL, 91.75 mmol) slowly over a period of 30 min. Checked for complete formation of mixed anhydride by NMR.

Added the compound of Formula VI (10 g, 61.3 mmol), DMAP (1.6 g, 13 mmol) and dry DMF (10 mL) and heated the mixture at reflux for about 7 h or until the reaction was complete (<3% compound III) by NMR. Cooled to room temperature, transferred the batch to a flask containing 2N H₂SO₄ (80 mL) slowly with agitation and continued agitation for about 30 min. Separated the layers and washed the organic layer with 5% NaHCO₃ (80 mL).

- 36 -


Concentrated the organic layer and crystallized the product from isopropyl alcohol (100 mL), filtered and dried. Yield: 20 g (92% molar); mp: 92-94°C.

Step (b):



Equipped a 3-necked 250 mL round bottom flask with a thermometer, an addition funnel and a nitrogen inlet. Added dry CH_2Cl_2 (20 mL) and neat borane dimethyl sulfide (2.82 mL, 28.2 mmol) and cooled the mixture to -5° to 0°C . Added a previously prepared toluene solution of (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo(1,2-c)(1,3,2) oxaza-borolidine (1.4 mL, 1.4 mmol, 5 mole %) to the mixture and stirred for 15 min. at $< 0^\circ\text{C}$. Added a solution of compound VII (10 g, 28.1 mmol) in CH_2Cl_2 (30 mL) slowly over a period of 3-4 h and maintained the reaction temperature between -5° to 0°C . Continued stirring for 1 to 2 h or until the reaction completed ($< 0.1\%$ compound VII) by NMR. Quenched the reaction by slowly adding CH_3OH (4 mL) while maintaining the temperature $< 0^\circ\text{C}$. Added 5% hydrogen peroxide (20 mL) followed by 4N H_2SO_4 (1.5 mL). Agitated the mixture for 15 min., separated the organic layer and washed with 2N H_2SO_4 (20 mL), 5% Na_2SO_3 (50 mL) and 10% NaCl (50 mL). Concentrated the organic layer to a low volume until water content is $< 0.05\%$. The product was used directly in the next step. Solution yield: $> 95\%$; de: 98%.


- 37 -

Step (c):


Equipped a 3-necked 500 mL round bottom flask with a thermometer, an addition funnel and a nitrogen inlet. Added the CH_2Cl_2 solution of compound VIII (10 g equivalent of compound VIII, 28.1 mmol) from step (b) and compound IX (12.05 g) and adjusted the total volume of the reaction mixture to 150 mL with dry CH_2Cl_2 . Cooled the mixture to -10°C and slowly added DIPEA (25.7 mL, 147.5 mmol) and maintained the temperature at $< -5^\circ\text{C}$. Added TMSCl (13.5 mL, 92.3 mmol) over a period of 30 min. while maintaining the reaction temperature $< -5^\circ\text{C}$. Agitated the reaction for 1 h or until the silylation was judged complete by NMR. Lowered the reaction temperature to -25 to -30°C . Added TiCl_4 (3.4 mL, 30.8 mmol) slowly and maintained the temperature $< -25^\circ\text{C}$. Agitated the reaction for 3 h at $< -25^\circ\text{C}$ and checked the reaction completion by NMR. Added glacial acetic acid (8 mL) slowly to the reaction mixture while maintaining the reaction temperature between -25 and -30°C . Poured the reaction mixture into 7% aqueous tartaric acid solution (140 mL) at 0°C , agitate for 1-2 h, and allowed the temperature to gradually rise to room temperature. Added 20% aqueous NaHSO_3 solution (50 mL) and continued agitation for another 2 h. Separated the organic layer and wash with water (120 mL). Concentrated the organic layer to a low volume and added bistrimethylsilyl acetamide (8.4 mL) and heated the mixture to reflux for 30 min. Concentrated the mixture to remove CH_2Cl_2 and crystallized the product from an ethyl acetate and heptane mixture, filtered, washed and dried to give 13 g (65% molar yield from compound VIII) of compound III. ^1H NMR (CDCl_3 , 400 MHz) δ 7.18(7H, m), 7.06(2H, t, $J=7.6\text{Hz}$), 6.98(2H, t, $J=8.66\text{Hz}$), 6.77(4H, m), 6.43(2H, dd, $J=8.84$, 4.4 Hz), 5.43(1H, dd, $J=8.13$, 2.21Hz), 4.47(1H, t, $J=8.57\text{ Hz}$), 4.46(2H, m), 4.29(1H, d, $J=9.51\text{Hz}$), 4.21(1H, dd, $J=8.71$, 2.91Hz), 1.55(3H, m), 1.41(1H, m), 0.28(9H, s),

- 38 -


-0.07(9H, s).

Step (d):**Example 1:**Intramolecular Cyclization of Compound III with Tetrabutylammonium Acetate:

Equipped a 1-necked 50 mL round bottom flask with a magnetic stirrer and a stopper. Added Compound III (1.0 g, 1.4 mmol), N,O-bis(trimethylsilyl) acetamide (1.0 mL), tetrabutylammonium acetate (0.020 g, 0.066 m mole or 5 mole%) and methyl t-butyl ether (10 mL). Stirred the mixture at room temperature for about 15 minutes. Analysis of the reaction mixture by ^1H NMR (CDCl_3 , 400 MHz) indicated the disappearance of compound III (specifically the disappearance of a characteristic doublet of a doublet at δ 6.43 ppm and appearance of a doublet of a triplet at 3.04 ppm ($J=7.51, 2.2\text{Hz}$) and a doublet at 4.54($J=2.2\text{ Hz}$). These two peaks are characteristic for C-3 and C-4 protons of a trans β -lactam respectively. HPLC analysis of an acidified sample of the reaction mixture also gave a peak at 21.2 min (96% area) matching the retention time of an authentic sample of compound XI.

Example 2:Intramolecular Cyclization of Compound III with Potassium Acetate and Tetrabutylammonium Bromide:


Equipped a 3-necked 125 mL round bottom flask with a magnetic stirrer. Added Compound III (5 g, 7.0 mmol), N,O-bis(trimethylsilyl) acetamide (5.0 mL), potassium acetate (0.035 g, 0.36m mole or 5 mole%), and tetrahydrofuran (50 mL). Stirred the mixture at room temperature for 0.5 h. Checked reaction progress by ¹HNMR which indicated no reaction whatsoever. Added tetrabutylammonium bromide (0.112 g, 0.35 mmol or 5 mole%). Stirred the mixture at room temperature for 0.7 h. Analysis of the reaction mixture by ¹H NMR was conducted in the same manner as for Example 1 above.

Example 3:Intramolecular Cyclization of Compound III with Cesium fluoride and tetrabutylammonium bromide:


Equipped a 3-necked 250 mL round bottom flask with a thermometer, an addition funnel and a nitrogen inlet. Added Compound III (5 g, 6.97 mmol), and anhydrous Tetrahydrofuran (50 mL). Stirred the mixture at room temperature for about 10 minutes, or until all solids dissolved. Added N,O Bistrimethylsilylacetamide

- 40 -

(BSA, 5 mL, 20.2 mmol). Stirred the solution at room temperature for about 15 minutes. Added Cesium Fluoride (CsF, 52.9 mg, 0.35 mmol, 5 mole %). Stirred the mixture at room temperature for about 45 minutes. Charged tetra-N-Butylammonium Bromide (TBABr, 112 mg, 0.35 mmol, 5 mole %). Stirred the mixture at room temperature for about 1 hour. Analysis of the reaction mixture by ^1H NMR was conducted in the same manner as for Example 1 above.

Example 4:Intramolecular Cyclization of Compound III with Tetrabutylammonium Hydroxide:

Equipped a 1-necked 50 mL round bottom flask with a magnetic stirrer and a stopper. Added compound III (1.0 g, 1.4 mmol), N,O-bis(trimethylsilyl) acetamide (1.0 mL), and tetrahydrofuran (10 mL). Stirred the mixture at room temperature for about 10 minutes. Added aqueous solution of tetrabutylammonium hydroxide (0.05mL; 40 w/w%; 0.077 mmole or 5.5 mole%). Stirred the mixture at room temperature for 1.3 h. Analysis by ^1H NMR showed that the reaction was only half way to completion. Added more N,O-bis(trimethylsilyl) acetamide (1.0 mL). Stirred the mixture at room temperature for 1 h. Analysis of the reaction mixture by ^1H NMR was conducted in the same manner as for Example 1 above.

Example 5:Intramolecular Cyclization of Compound III with Potassium Acetate:

Equipped a 1-necked 50 mL round bottom flask with a magnetic stirrer and a stopper. Added compound III (1.0 g, 1.4 mmol), N,O-bis(trimethylsilyl) acetamide (1.0 mL), potassium acetate (0.14 g, 1.4 mmol), and tetrahydrofuran (10 mL). Stirred the mixture at room temperature for about 1.5 h. Analysis by ¹H NMR showed no reaction. Heated the mixture to about 50°C, and stirred the mixture at that temperature for about 3 h. Analysis by ¹H NMR showed that the reaction was only about 65% complete. Again, heated the mixture to about 60 °C, and stirred the mixture at that temperature for about 1 h. Analysis of the reaction mixture by ¹H NMR was conducted in the same manner as for Example 1 above.

Step (e):

Added a premixed solution of isopropyl alcohol and 2N H₂SO₄ to the product of step (d) and agitated the mixture at room temperature for 1 h. Crystallized compound IV from aqueous isopropyl alcohol. Filtered the product and washed with dilute aqueous isopropyl alcohol followed by water until the pH of wash was <5. Dried the product at 60°C in a draft oven or under vacuum to give compound IV.

It will be appreciated by those skilled in the art that changes could be made to the embodiments described above without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications that are within the spirit and scope of the invention, as defined by the appended claims.

- 42 -

THEREFORE, WE CLAIM:

1. A process for preparing an azetidinone, comprising the step of reacting:

(1) a β -(substituted-amino)amide, a β -(substituted-amino)-acid ester, or a β -(substituted-amino)thiolcarbonic acid ester with

(2) at least one silylating agent and

(3) at least one cyclizing agent which is

(a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof, or


(b) the reaction product of:

(i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or

(ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride,

wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl.

2. The process according to claim 1, wherein the azetidinone is represented by the structural Formula (I):



or pharmaceutically acceptable salts or solvates of the azetidinone of Formula (I), wherein in Formula (I) above:

- 43 -

X', Y' and Z can be the same or different and each is independently selected from the group consisting of -CH₂-, -CH(alkyl)- and -C(alkyl)₂-;

Q¹ and Q² can be the same or different and each is independently selected from the group consisting of H, -(C₀-C₃₀ alkylene)-G, -OR⁶, -OC(O)R⁶, -OC(O)OR⁹,
 5 and -OC(O)NR⁶R⁷M;

Q³, Q⁴, and Q⁵ can be the same or different and each is independently 1 to 5 substituents independently selected from the group consisting of acyl, alkyl, alkylaryl, alkylheteroaryl, alkylsulfonyl alkenyl, alkoxy, alkoxy carbonyl, alkynyl,, -(C₀-C₃₀ alkylene)-G,

10 -(C₀-C₁₀ alkylene)-OR⁶, -(C₀-C₁₀ alkylene)-C(O)R⁶,
 -(C₀-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-OC(O)R⁶,
 -(C₀-C₁₀ alkylene)-OC(O)OR⁹, -CH=CH-C(O)R⁶, -CH=CH-C(O)OR⁶,
 -C≡C-C(O)OR⁶, -C≡C-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-OR⁶,
 -O-(C₁-C₁₀ alkylene)-C(O)R⁶, -O-(C₁-C₁₀ alkylene)-C(O)OR⁶, -CN,
 15 -C(=N-CN)-NH₂, -C(=NH)-NHR¹⁰, -O-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -O-(C₀-C₁₀ alkylene)-C(O)NR⁶NR⁷C(O)OR⁶,
 -O-(C₁-C₁₀ alkylene)-C(O)(aryl)-N-N=N⁻,
 -OC(O)-(C₁-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -(C₀-C₁₀ alkylene)-OC(O)NR⁶R⁷, -NO₂, -(C₀-C₁₀ alkylene)-NR⁶R⁷,
 20 -O-(C₂-C₁₀ alkylene)-NR⁶R⁷, -NR⁶C(O)R⁷, -NR⁶C(O)OR⁹, -NR⁶C(O)NR⁷R⁸,
 -NR⁶S(O)₀₋₂R⁹, -N(S(O)₀₋₂R⁹)₂, -CHNOR⁶, -C(O)NR⁶R⁷, -C(O)NR⁶NR⁶R⁷,
 -S(O)₀₋₂NR⁶R⁷, -S(O)₀₋₂R⁹, -O-C(O)-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,
 -OC(O)-(C₁-C₁₀ alkylene)-NR⁶C(O)O-(alkylaryl), -P(O)(OR¹⁰)₂,
 -(C₁-C₁₀ alkylene)-OSi(alkyl)₃, -CF₃, -OCF₃, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,
 25 alkoxy carbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidioyl, allyl, allyloxy,
 aryloxy carbonyl, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, arylsulfonyl,
 aroylaroyloxy, aroyl, arylalkoxy carbonyl, benzoylbenzoyloxy, carboxy, cyano,
 cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl,
 heteroarylalkenyl, heteroarylalkynyl, heteroarylalkyl, heteroarylalkoxy,

- 44 -

heteroarylsulfonyl, heteroarylthio, dioxolanyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy, hydroxy, hydroxyalkyl, and alkylsulfonyl;

wherein optionally one or more carbon atoms of the $-(C_0-C_{30} \text{ alkylene})-$ radical of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is independently replaced by $-O-$, $-C(O)-$, $-CH=CH-$, $-C\equiv C-$, $-N(\text{alkyl})-$, $-N(\text{alkylaryl})-$ or $-NH-$;

G is selected from the group consisting of a sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue, oligopeptide residue comprising 2 to 9 amino acids, trialkylammoniumalkyl radical and $-S(O)_2-OH$,

R^2 and R^3 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl and aryl;

R^6 , R^7 and R^8 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, arylalkyl and a protecting group; and

each R^9 is independently alkyl, cycloalkyl, aryl or arylalkyl.

each R^{10} is independently H or alkyl;


q is 0 or 1;

r is 0 or 1;

m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5.


3. The process according to claim 2, wherein the azetidinone is represented by the structural Formula (IV):

- 45 -



(IV).

4. The process according to claim 1, wherein the β -(substituted-amino)amide comprises a carbamoyl portion B-C(O)-, wherein B is a deprotonated chiral auxiliary selected from the group consisting of



wherein X is -O-, -S- or -N(alkyl)-; Y is =O or =S; and R¹² and R¹³ are each independently selected from the group consisting of alkyl, aryl, arylalkyl, and alkoxy, or wherein one of R¹² or R¹³ is as defined above and the other is hydrogen.


5. The process according to claim 4, wherein the aryl group of R¹² or R¹³ is independently selected from the group consisting of phenyl, naphthyl, benzyl, substituted phenyl, substituted naphthyl and substituted benzyl, wherein the substituents on the phenyl or naphthyl are 1-3 substituents selected from the group consisting of alkyl, alkoxy, phenyl and benzyl.

6. The process according claim to 4, wherein the β -(substituted-amino)amide comprises a carbamoyl portion B-C(O)-, where B is (R¹⁴)(R¹⁵)N-, and

- 46 -

R¹⁴ and R¹⁵ are each independently selected from the group consisting of alkyl, aryl, arylalkyl.

7. The process according to claim 4, wherein the β -(substituted-amino)amide is represented by Formula (II):



wherein B is as defined in claim 4,

X', Y' and Z can be the same or different and each is independently selected from the group consisting of -CH₂-, -CH(alkyl)- and -C(alkyl)₂-;

Q¹ and Q² can be the same or different and each is independently selected from the group consisting of H, -(C₀-C₃₀ alkylene)-G, -OR⁶, -OC(O)R⁶, -OC(O)OR⁹, and -OC(O)NR⁶R⁷;

Q³, Q⁴, and Q⁵ can be the same or different and each is independently 1 to 5 substituents independently selected from the group consisting of acyl, alkyl, alkylaryl, alkylheteroaryl, alkylsulfonyl alkenyl, alkoxy, alkoxy carbonyl, alkynyl, -(C₀-C₃₀ alkylene)-G,

-(C₀-C₁₀ alkylene)-OR⁶, -(C₀-C₁₀ alkylene)-C(O)R⁶,

-(C₀-C₁₀ alkylene)-C(O)OR⁶, -(C₀-C₁₀ alkylene)-OC(O)R⁶,

- 47 -

$-(C_0-C_{10} \text{ alkylene})-OC(O)OR^9$, $-CH=CH-C(O)R^6$, $-CH=CH-C(O)OR^6$,
 $-C\equiv C-C(O)OR^6$, $-C\equiv C-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-OR^6$,
 $-O-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-CN$,
 $-C(=N-CN)-NH_2$, $-C(=NH)-NHR^{10}$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,
 $-O-(C_0-C_{10} \text{ alkylene})-C(O)NR^6NR^7C(O)OR^6$,
 $-O-(C_1-C_{10} \text{ alkylene})-C(O)(\text{aryl})-N=N=N^+$,
 $-OC(O)-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-(C_0-C_{10} \text{ alkylene})-C(O)NR^6R^7$,
 $-(C_0-C_{10} \text{ alkylene})-OC(O)NR^6R^7$, $-NO_2$, $-(C_0-C_{10} \text{ alkylene})-NR^6R^7$,
 $-O-(C_2-C_{10} \text{ alkylene})-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$,
 $-NR^6S(O)_{0-2}R^9$, $-N(S(O)_{0-2}R^9)_2$, $-CHNOR^6$, $-C(O)NR^6R^7$, $-C(O)NR^6NR^6R^7$,
 $-S(O)_{0-2}NR^6R^7$, $-S(O)_{0-2}R^9$, $-O-C(O)-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,
 $-OC(O)-(C_1-C_{10} \text{ alkylene})-NR^6C(O)O-(\text{alkylaryl})$, $-P(O)(OR^{10})_2$,
 $-(C_1-C_{10} \text{ alkylene})-OSi(\text{alkyl})_3$, $-CF_3$, $-OCF_3$, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy,
alkoxycarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkylidioyl, allyl, allyloxy,
aryloxycarbonyl, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, arylsulfonyl,
aroylaroyloxy, aroyl, arylalkoxycarbonyl, benzoylbenzoyloxy, carboxy, cyano,
cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl,
heteroarylalkenyl, heteroarylalkynyl, heteroarylalkyl, heteroarylalkoxy,
heteroarylsulfonyl, heteroarylthio, dioxolanyl, heterocyclyl, heterocyclylalkyl,
heterocyclylcarbonyl, heterocyclylcarbonylalkoxy, hydroxy, hydroxyalkyl, and
alkylsulfonyl;

wherein optionally one or more carbon atoms of the $-(C_0-C_{30} \text{ alkylene})$ - radical of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is independently replaced by

$-O-$, $-C(O)-$, $-CH=CH-$, $-C\equiv C-$, $-N(\text{alkyl})-$, $-N(\text{alkylaryl})-$ or $-NH-$;

G is selected from the group consisting of a sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue, oligopeptide residue comprising 2 to 9 amino acids, trialkylammoniumalkyl radical and $-S(O)_2-OH$,

R^2 and R^3 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl and aryl;

- 48 -

R^6 , R^7 and R^8 can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, arylalkyl and a protecting group; and

each R^9 is independently alkyl, cycloalkyl, aryl or arylalkyl.


each R^{10} is independently H or alkyl;

q is 0 or 1;

r is 0 or 1;

m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5.

8. The process according to claim 7, wherein the β -(substituted-amino)amide is represented by Formula (III):



(III).

9. The process according claim 1, wherein the β -(substituted-amino)acid ester comprises a carboxylic acid ester portion R^{14} -O-C(O)-, wherein R^{14} is alkyl or aryl.

- 49 -

10. The process according claim 1, wherein the β -(substituted-amino)thiolcarbonic acid ester comprises a thiolcarbonic acid ester portion R^{14} -S-C(O)-, wherein R^{14} is alkyl or aryl.

5 11. The process according claim 1, wherein the silylating agent is a silylenol ether.

12. The process according claim 11, wherein the silylating agent is selected from the group consisting of bistrimethylsilylacetamide, N-methyl-O-trimethyl
10 silylacetamide or isopropenyloxy trimethylsilane.

13. The process according to claim 1, wherein the alkali metal carboxylate is a carboxylate salt of an alkali metal selected from the group consisting of lithium,
15 sodium, potassium and cesium.

14. The process according to claim 1, wherein the alkali metal carboxylate or quaternary ammonium carboxylate comprises a carboxylate moiety selected from the group consisting of formate, acetate, propionate, butyrate, valerate, caproate,
20 caprylate, laurate, myristate, palmitate, stearate, oleate, linoleate, linolenate, cyclohexanecarboxylate, phenylacetate, benzoate and toluate.

15. The process according to claim 14, wherein the carboxylate moiety of the alkali metal carboxylate is acetate.
25

16. The process according to claim 14, wherein the alkali metal carboxylate is potassium acetate.

17. The process according to claim 14, wherein the carboxylate moiety of the quaternary ammonium carboxylate is acetate.
30

- 50 -

18. The process according to claim 1, wherein the quaternary ammonium moiety of the cyclizing agent is tetra n-butylammonium.

19. The process according to claim 14, wherein the quaternary ammonium
5 carboxylate is tetra n-butylammonium acetate.

20. The process according to claim 1, wherein the quaternary ammonium hydroxide is tetra n-butylammonium hydroxide.

10 21. The process according to claim 1, wherein the quaternary ammonium halide comprises a halide moiety selected from the group consisting of fluoride, chloride, bromide and iodide.

22. The process according to claim 1, wherein the cyclizing agent is the
15 reaction product of tetra n-butylammonium bromide and potassium acetate.

23. The process according to claim 1, wherein the cyclizing agent is the reaction product of tetra n-butylammonium bromide and cesium fluoride.

20 24. The process according to claim 1, wherein the reaction product (3)(b) is formed in situ in the presence of reactants (1) and (2).

25 25. The process according to claim 1, further comprising the step of forming the reaction product (3)(b) prior to reaction with reactants (1) and (2).


26. The process according to claim 1, wherein the silylating agent is bistrimethylsilylacetamide and the cyclizing agent is tetra n-butylammonium acetate.

30 27. The process according to claim 1, further comprising adding a second cyclizing agent which is a fluoride ion catalyst.

- 51 -

28. The process according to claim 1, comprising pre-reacting reactant (1) with reactant (2).


29. The process according to claim 1, for preparing an azetidinone represented by the structural Formula (IV):



(IV)

comprising reacting:

(1) a β -(substituted-amino)amide of the formula (III)



(III)

with (2) at least one silylating agent and
(3) at least one cyclizing agent which is


(a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof; or

(b) the reaction product of:

- 52 -


- (i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or
- (ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride,
- wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl.

30. A process for preparing a compound represented by the Formula (IV):



comprising the step of:

cyclizing the β -(substituted-amino)amide of formula XII



wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxycarbonyl, and wherein Prot is a silyl protecting group

with at least one silylating agent and at least one cyclizing agent which is


- (a) selected from the group consisting of alkali metal carboxylates, quaternary ammonium carboxylates, quaternary ammonium hydroxides, quaternary ammonium alkoxides, quaternary ammonium aryloxides and hydrates thereof; or
- (b) the reaction product of:

(i) at least one quaternary ammonium halide and at least one alkali metal carboxylate; or

(ii) at least one quaternary ammonium chloride, quaternary ammonium bromide, or quaternary ammonium iodide and at least one alkali metal fluoride,

wherein a quaternary ammonium moiety of the cyclizing agent is unsubstituted or substituted by one to four groups independently selected from the group consisting of alkyl, arylalkyl and arylalkyl-alkyl,


10 to obtain the compound of Formula XI:



removing the silyl protecting groups to form the compound of Formula (IV).

31. The process according to claim 30, further comprising the steps of:

reacting a chiral alcohol of Formula VIII, an imine of Formula IX and a silyl protecting agent, then condensing the silyl-protected compounds to obtain a β -(substituted-amino)amide of Formula XII, wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxycarbonyl, and wherein Prot is a silyl protecting group:




prior to cyclizing the β -(substituted-amino)amide of Formula XII.

32. The process according to claim 30, further comprising the steps of:


- 54 -

(a) reacting p-fluorobenzoylbutyric acid of Formula V with pivaloyl chloride and acylating the product with a chiral auxiliary of Formula VI to obtain a ketone of Formula VII:




wherein X is -O-, -S- or -N(C₁-C₆ alkyl); Y is =O or =S; and R¹ is alkyl, aryl or alkoxy carbonyl;

(b) reducing the ketone of Formula VII in the presence of a chiral catalyst to an alcohol of Formula VIII:



(c) reacting the chiral alcohol of Formula VIII, an imine of formula IX and a silyl protecting agent, then condensing the silyl-protected compounds to obtain a β-(substituted-amino)amide of Formula XII, wherein Prot is a silyl protecting group:



prior to cyclizing the β-(substituted-amino)amide of Formula XII.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/045901

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D205/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6 093 812 A (THIRUVENGADAM ET AL) 25 July 2000 (2000-07-25) column 13	1-32
Y	WO 00/34240 A (SCHERING CORPORATION) 15 June 2000 (2000-06-15) page 10	1-32
Y	EP 0 707 567 A (SCHERING CORPORATION) 24 April 1996 (1996-04-24) cited in the application claims 8,9	1-32
Y	US 6 207 822 B1 (THIRUVENGADAM TIRUVETTIPURAM K ET AL) 27 March 2001 (2001-03-27) cited in the application column 7	1-32

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

11 May 2006

Date of mailing of the international search report

23/05/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2005/045901

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6093812	A	25-07-2000	NONE	
WO 0034240	A	15-06-2000	AT 297892 T	15-07-2005
			AU 2030100 A	26-06-2000
			CA 2353981 A1	15-06-2000
			CN 1130342 C	10-12-2003
			DE 69925862 D1	21-07-2005
			DE 69925862 T2	04-05-2006
			EP 1137634 A1	04-10-2001
			ES 2244238 T3	01-12-2005
			HK 1039487 A1	12-08-2005
			JP 3640888 B2	20-04-2005
			JP 2002531546 T	24-09-2002
			JP 2005053931 A	03-03-2005
			PT 1137634 T	31-10-2005
			ZA 200104004 A	16-08-2002
EP 0707567	A	24-04-1996	AT 205475 T	15-09-2001
			AU 681419 B2	28-08-1997
			AU 7214694 A	06-02-1995
			CA 2166784 A1	19-01-1995
			CN 1126989 A	17-07-1996
			CZ 9600076 A3	17-04-1996
			DE 69428280 D1	18-10-2001
			DE 69428280 T2	13-06-2002
			DK 707567 T3	26-11-2001
			ES 2160125 T3	01-11-2001
			FI 960095 A	09-01-1996
			HU 73237 A2	29-07-1996
			JP 2840452 B2	24-12-1998
			JP 8507085 T	30-07-1996
			KR 176001 B1	20-03-1999
			MX 9405152 A1	31-01-1995
			NO 960083 A	04-03-1996
			NZ 268798 A	29-01-1997
			PL 312502 A1	29-04-1996
			PT 707567 T	30-01-2002
			SK 3796 A3	05-02-1997
			WO 9501961 A1	19-01-1995
			US 5728827 A	17-03-1998
US 6207822	B1	27-03-2001	NONE	